

COMPUTING THE CENTER OF AREA OF A CONVEX POLYGON*

Peter Braß[†]

Laura Heinrich-Litan[‡]

Pat Morin[§]

ABSTRACT. The center of area of a convex planar set X is the point p for which the minimum area of X intersected by any halfplane containing p is maximized. We describe a simple randomized linear-time algorithm for computing the center of area of a convex n -gon.

1 Introduction

Let X be a convex planar set with unit area. A *center of area* of X is a point p^* that maximizes the *cut off area function*

$$f(p) = \min\{\text{area}(h \cap X) \mid h \text{ is a halfplane that contains } p\} ,$$

and the value $\delta^* = f(p^*)$ is known as *Winternitz's measure of symmetry* [14]. The δ -level Γ_δ of X is defined as

$$\Gamma_\delta = \{p \mid f(p) = \delta\} .$$

It is known that Γ_δ is a simple closed convex curve and that Γ_{δ_1} is strictly contained in Γ_{δ_2} if $\delta_1 > \delta_2$. From this it follows that p^* is unique.

There is a long history of work on the center of area of convex sets. A classical result of Winternitz [3, pp. 54–55], which has been rediscovered many times [12, 16–19], is that $f(g) \geq 4/9$ where g is the centroid of X , with equality if and only if X is a triangle. (In d dimensions, Ehrhart [11] showed that $f(g) \geq d^d/(d+1)^d$ with equality if and only if X is a pyramid on any $(d-1)$ -dimensional convex base.) For centrally symmetric sets, $f(g) = 1/2$, since any line through the point of symmetry cuts X into two pieces of equal area. Thus, $4/9 \leq f(g) \leq 1/2$ with $f(g) = 4/9$ for triangles and $f(g)$ close to $1/2$ for highly symmetric sets.

Although much is known about the center of area, it is quite nontrivial to determine the center of area for a given convex set. In a series of papers, Díaz and O'Rourke [7, 8, 9] develop an $O(n^6 \log^2 n)$ time algorithm for finding the center of area of a convex n -gon. The same authors give an approximation algorithm that runs in $O(GK(n+K))$ time, where G is the bit-precision of the input polygon P and K is the output bit-precision of the point p^* . Braß and Heinrich-Litan [4] describe an $O(n^2 \log^3 n \alpha(n))$ time algorithm for computing the center of area of a convex n -gon. As an application of tools for searching in arrangements of lines, Langerman and Steiger [15] present an $O(n \log^3 n)$ time algorithm for finding the center of area of a convex n -gon. All of these algorithms are deterministic.

In this paper we give a simple randomized linear-time algorithm for finding the center of area of a convex n -gon P , which also computes Winternitz's measure of symmetry for P . We proceed by first giving a linear-time algorithm for the decision problem: Does there exist a point p such that $f(p) > \delta$? We then

* This research was partly funded by Deutsche Forschungsgemeinschaft (DFG), grant BR 1465/5-2, and by the IST Programme of the EU as a Shared-cost RTD (FET Open) Project under Contract Number IST-2000-26473 (ECG - Effective Computational Geometry for Curves and Surfaces).

[†]Department of Computer Science R8/206, City College CUNY, 138th Street at Convent Ave., New York NY 10031, USA, peter@cs.ccny.cuny.edu

[‡]Institut für Informatik, Freie Universität Berlin, D-14195 Berlin, Germany, litan@inf.fu-berlin.de

[§]School of Computer Science, Carleton University, 1125 Colonel By Drive, Ottawa, Ontario, K1S 5B6, Canada, morin@cs.carleton.ca

apply a randomized technique due to Chan [5] to turn this decision algorithm into a linear-time optimization algorithm. For convenience, our model of computation is the real-RAM, though we do not use any functions that are specific to this model. We require only that it is possible to compute the exact area of a convex polygon.

The remainder of the paper is organized as follows. Section 2 describes our algorithm for the decision problem and Section 3 shows how to convert this decision algorithm into an optimization algorithm. Section 4 summarizes and concludes with directions for future research.

2 The Decision Algorithm

In this section, we give an $O(n)$ time algorithm for the following decision problem: Is there a point p such that $f(p) \geq \delta$? An alternative statement of this problem is: is Γ_δ non-empty? In what follows, we show that Γ_δ can be computed in $O(n)$ time.

A δ -cut of P is a directed line segment uv with endpoints u and v on the boundary of P such that the area of P to the right of uv is at most δ . Note that, for any δ -cut uv , the point p can not be to the right of uv . On the other hand, if there is no δ -cut uv with p on its right, then $f(p) \geq \delta$. Therefore, each δ -cut defines a linear constraint on the location of p , which we call a δ -constraint. The answer to the decision problem is affirmative if and only if there is a point p that simultaneously satisfies all δ -constraints. If such a point p exists, we call the constraints feasible, otherwise we call them infeasible.

Unfortunately, every polygon has an infinite number of δ -cuts and hence an infinite number of δ -constraints. However, we will show that all constraints imposed by these δ -cuts can be represented succinctly as $O(n)$ non-linear (but convex) constraints that can be computed in $O(n)$ time.

To generate a representation of all δ -constraints, we begin by choosing a point u on the boundary of P and finding the unique point v so that uv is a δ -cut. Next, we sweep the points u and v counterclockwise along the boundary of P maintaining the invariant that uv has an area of exactly δ to its right. We continue this process until we return to the original points u and v .

Observe that, as long as u and v do not cross a vertex of P , the intersection of all δ -constraints belonging to an edge pair is a convex region whose boundary consists of at most 2 linear pieces and 1 non-linear piece. (See Figure 1.) In fact, this non-linear piece is a hyperbolic arc, since it consists of midpoints of δ -cuts. Furthermore, the description complexity of these pieces is constant, since they are defined by four vertices of P and one area. Thus, the intersection of all these δ -constraints can be computed explicitly in constant time. Since u and v sweep over each vertex exactly once, we obtain $2n$ such convex constraints whose intersection is equal to the intersection of all δ -constraints.

Therefore, the decision problem reduces to determining if the intersection of $2n$ convex regions is empty. We can compute an explicit representation of this intersection in $O(n)$ time, as follows: Separately compute the intersection of all δ -constraints that contain the point $(0, +\infty)$ and all δ -constraints that contain the point $(0, -\infty)$ and then compute the intersection of the two resulting convex regions. Since the δ -constraints are generated sorted by slope, the first step is easily done in $O(n)$ time using an algorithm similar to Graham's Scan [1, 13]. Since the two boundaries of the two resulting regions are x -monotone and upwards, respectively downwards, convex, their intersection (step two) can be computed in $O(n)$ time using a left-to-right plane sweep [2].

We have just proven:

Theorem 1. *There exists an $O(n)$ time algorithm for the decision problem: Does there exist a point p such that $f(p) \geq \delta$?*




Figure 1: The intersection of all δ -constraints defined by a pair of edges makes a convex region whose boundary consists of at most 3 pieces.

3 The Optimization Algorithm

In this section, we show how to use the decision algorithm of the previous section along with a technique of Chan [5] to solve the optimization problem: What is the largest value of δ such that Γ_δ is non-empty? Chan's technique requires only that we be able to (1) solve the decision problem in $D(n) = \Omega(n^\epsilon)$ time, $\epsilon > 0$ and (2) generate a set of r subproblems each of size αn , $\alpha < 1$, such that the solution to the original problem is the minimum (or maximum) of the solutions to the subproblems. Under these conditions, the optimization problem can be solved by a randomized algorithm in $O(D(n))$ expected time.

To apply Chan's technique, we need a suitable definition of subproblem. Let S be a subset of edges of P . The S -induced δ -constraints are the set of all δ -constraints uv , where u and v are both on edges of S . The type of subproblems we consider are those of determining for a given set S and a value δ whether or not the S -induced δ -constraints are feasible. To obtain a linear-time algorithm, we must be able to solve such subproblems in $O(|S|)$ time.

For a given set S , computing a representation of the S -induced δ -constraints, can be done using a modification of the sweep algorithm from the previous section so that it only considers δ -cuts uv where u and v are on elements of S . The only technical tool required for this modification is a data structure that, given two points u and v on elements of S (the boundary of P) tells us the area of P to the right of uv in constant time. This data structure is provided by Czyzowicz *et al* [6] who show that any convex n -gon can be preprocessed in $O(n)$ time so that the area of the polygon to the right of any chord uv can be computed in $O(1)$ time. Using this data structure, it is straightforward to generate a representation of S -induced δ -constraints in $O(|S|)$ time. Once we have computed these constraints, we can test if they are feasible in $O(|S|)$ time. Thus, Condition 1 required to use Chan's technique is satisfied with $D(n) = \Theta(n)$.

Next, we observe that Helly's theorem in the plane (c.f., Eckhoff [10]) implies that for any $\delta > \delta^*$ there exists a set of three δ -constraints whose intersection is empty. In our context, this means that P contains 6 edges such that, for any $\delta > \delta^*$, the δ -constraints induced by those edges are infeasible. Therefore, if a set S contains those 6 edges, then the S -induced δ -constraints are feasible if and only if $\delta \leq \delta^*$.

Therefore, we can solve our maximization problem as follows: Partition the edges of P in 7 groups, E_1, \dots, E_7 , each of size approximately $n/7$. Next, generate subsets S_1, \dots, S_7 , by taking all 7 6-tuples of E_1, \dots, E_7 . Note that, for each S_i , the S_i -induced δ -constraints are satisfiable if $\delta \leq \delta^*$, since they are just a subset of the original constraints. On the other hand, for the set S_j that contains the 6 edges guaranteed by Helly's theorem, the S_j -induced δ -constraints are not satisfiable for any $\delta > \delta^*$. Therefore,

$$\delta^* = \min \{ \max \{ \delta \mid S_i\text{-induced } \delta\text{-constraints are satisfiable} \} \mid 1 \leq i \leq 7 \} .$$

Finally, observe that each S_i is of size at most αn , for $\alpha = 6/7$. Therefore, we have satisfied the second condition required to apply Chan's optimization technique. This completes the proof of:

Theorem 2. *There exists a randomized $O(n)$ expected time algorithm for the optimization problem: What is the largest value δ^* for which Γ_{δ^*} is non-empty?*

Of course, once δ^* is known, an explicit representation of Γ_{δ^*} can be computed in $O(n)$ time. Alternatively, Chan's technique can also be made to output a point $p^* \in \Gamma_{\delta^*}$ [5].

4 Conclusions

We have given a randomized linear-time algorithm for determining the center of area of a convex n -gon. This algorithm is simple, implementable, and is asymptotically faster than any previously known algorithm.

Although our algorithm is simple and easy to implement, the constants are very large. A close examination of Chan's technique reveals that the number of subproblems generated in our application is actually $r \geq \binom{k}{6}$, where k is an integer that satisfies $\ln \binom{k}{6} + 1 < k/6$. The smallest such value of k is 146, which leads to $r = \binom{146}{6} = 12\,122\,560\,164$ subproblems. Reducing this constant while maintaining the $O(n)$ asymptotic running time remains an open problem. A linear-time deterministic algorithm is also an open problem.

Finally, we have not considered the problem of computing the center of area of a non-convex polygon. There are two different versions of this problem, depending on whether a cut is defined as a chord of P , which partitions P into two polygons, or a line which may partition P into many polygons. Approximation algorithms for the second case are considered by Díaz and O'Rourke [7]. To the best of our knowledge, there are no exact algorithms for either version.

Acknowledgements

The authors would like to thank Stefan Langerman for helpful discussions and for bringing his paper [15] to our attention.

References

- [1] A. M. Andrew. Another efficient algorithm for convex hulls in two dimensions. *Information Processing Letters*, 9(5):216–219, 1979.
- [2] J. L. Bentley and T. A. Ottmann. Algorithms for reporting and counting geometric intersections. *IEEE Transactions on Computers*, C-28:643–647, 1979.
- [3] W. Blaschke. *Vorlesungen über Differentialgeometrie. II, Affine Differentialgeometrie*. Springer, Berlin, 1923.
- [4] P. Braß and L. Heinrich-Litan. Computing the center of area of a convex polygon. Technical Report B 02-10, Freie Universität Berlin, Fachbereich Mathematik und Informatik, March 2002.
- [5] T. Chan. Geometric applications of a randomized optimization technique. *Discrete & Computational Geometry*, 22:547–567, 1999.
- [6] J. Czyzowicz, F. Contreras-Alcalá, and J. Urrutia. On measuring areas of polygons. In *Proceedings of the 10th Canadian Conference on Computational Geometry (CCCG'98)*, 1998. URL <http://www.cccg.ca/proceedings/1998/>.

- [7] M. Díaz and J. O'Rourke. Computing the center of area of a polygon. In F. Dehne, J.-R. Sack, and N. Santoro, editors, *Workshop on Algorithms and Data Structures (WADS '89)*, volume 382 of *Lecture Notes in Computer Science*, pages 171–182, Ottawa, Canada, 1989. Springer-Verlag.
- [8] M. Díaz and J. O'Rourke. Chord centers for convex polygons. *Contemporary Mathematics*, 119:29–43, 1991. Proceedings of the AMS Special Session on Vision Geometry 1989.
- [9] M. Díaz and J. O'Rourke. Algorithms for computing the center of area of a convex polygon. *The Visual Computer*, 10:432–442, 1994.
- [10] J. Eckhoff. Helly, Radon, and Caratheodory type theorems. In P. M. Gruber and J. M. Wills, editors, *Handbook of Convex Geometry*, chapter 2.1, pages 389–448. North-Holland, Amsterdam, Netherlands, 1993.
- [11] E. Ehrhart. Sur les ovales and les ovoïdes. *C. R. Acad. Sci. Paris*, 240:583–585, 1955.
- [12] E. Ehrhart. Une généralisation du théorème de Minkowski. *C. R. Acad. Sci. Paris*, 240:483–485, 1955.
- [13] R. L. Graham. An efficient algorithm for determining the convex hull of a finite planar set. *Information Processing Letters*, 1(4):132–133, June 1972.
- [14] B. Grünbaum. Measures of symmetry for convex sets. In V. Klee, editor, *Convexity: AMS Proceedings of the Symposium on Pure Mathematics VII*, pages 233–270, 1963.
- [15] S. Langerman and W. Steiger. Optimization in arrangements. Technical Report SOCS-02.7, School of Computer Science, McGill University, 2002.
- [16] M. A. Lavrent'ev and L. A. Lyusternik. *Elements of the Calculus of Variations*, volume I, part II. Moscow, 1935. (in Russian).
- [17] B. H. Neumann. On an invariant of plane regions and mass distributions. *Journal of the London Mathematical Society*, pages 226–237, 1945.
- [18] D. J. Newman. Partitioning of areas by straight lines. *Notices of the American Mathematical Society*, 5: 510, 1958.
- [19] I. M. Yaglom and V. G. Boltyanskii. *Convex Figures*. Holt, Rinehart and Winston, 1961. (Translated from Russian).